

- American Thoracic Society. Management of malignant pleural effusions. *Am J Respir Crit Care Med* 2000;162:1987–2001.
- Antunes G, Neville E. Management of malignant pleural effusions. *Thorax* 2000;55:981–983.
- Roberts ME, Neville E, Berrisford RG, et al. Management of a malignant pleural effusion: British Thoracic Society Pleural Disease Guideline 2010. *Thorax* 2010;65:ii32–ii40.
- Feller-Kopman D, Berkowitz D, Boiselle P, et al. Large-volume thoracentesis and the risk of reexpansion pulmonary edema. *Ann Thorac Surg* 2007;84:1656–1661.
- Sahn SA. Pleural effusion in lung cancer. *Clin Chest Med* 1993;14:189–200.
- DeCamp MM Jr, Jaklitsch MT, Mentzer SJ, et al. The safety and versatility of video-thoracoscopy: a prospective analysis of 895 consecutive cases. *J Am Coll Surg* 1995;181:113–120.
- Davies CW, Traill ZC, Gleeson FV, et al. Intrapleural streptokinase in the management of malignant multiloculated pleural effusions. *Chest* 1999;115:729–733.
- Kessinger A, Wigton RS. Intracavitary bleomycin and tetracycline in the management of malignant pleural effusions: a randomized study. *J Surg Oncol* 1987;36:81–83.
- Mager HJ, Maesen B, Verzijlbergen F, et al. Distribution of talc suspension during treatment of malignant pleural effusion with talc pleurodesis. *Lung Cancer* 2002;36:77–81.
- Robinson LA, Fleming WH, Galbraith TA. Intrapleural doxycycline control of malignant pleural effusions. *Ann Thorac Surg* 1993;55:1115–1121.
- Patz EF Jr, McAdams HP, Erasmus JJ, et al. Sclerotherapy for malignant pleural effusions: a prospective randomized trial of bleomycin vs doxycycline with small-bore catheter drainage. *Chest* 1998;113:1305–1311.
- Lee YC, Baumann MH, Maskell NA, et al. Pleurodesis practice for malignant pleural effusions in five English-speaking countries: survey of pulmonologists. *Chest* 2003;124:2229–2238.
- Heffner JE, Standerfer RJ, Torstveit J, et al. Clinical efficacy of doxycycline for pleurodesis. *Chest* 1994;105:1743–1747.
- Hamed H, Fentiman IS, Chaudary MA, et al. Comparison of intracavitary bleomycin and talc for control of pleural effusions secondary to carcinoma of the breast. *Br J Surg* 1989;76:1266–1267.
- Colt HG, Russack V, Chiu Y, et al. A comparison of thoracoscopic talc insufflation, slurry, and mechanical abrasion pleurodesis. *Chest* 1997;111:442–448.
- Fentiman IS, Rubens RD, Hayward JL. A comparison of intracavitary talc and tetracycline for the control of pleural effusions secondary to breast cancer. *Eur J Cancer Clin Oncol* 1986;22:1079–1081.
- Hartman DL, Gaitner JM, Kesler KA, et al. Comparison of insufflated talc under thoracoscopic guidance with standard tetracycline and bleomycin pleurodesis for control of malignant pleural effusions. *J Thorac Cardiovasc Surg* 1993;105:743–747.
- Viallat JR, Rey F, Astoul P, et al. Thoracoscopic talc poudrage pleurodesis for malignant effusions: a review of 360 cases. *Chest* 1996;110:1387–1393.
- Dresler CM, Olak J, Herndon JE, et al. Phase III intergroup study of talc poudrage vs talc slurry sclerosis for malignant pleural effusion. *Chest* 2005;127:909–915.
- Stefani A, Natali P, Casali C, et al. Talc poudrage versus talc slurry in the treatment of malignant pleural effusion. A prospective comparative study. *Eur J Cardiothorac Surg* 2006;30:827–832.
- Tan C, Sedrakyan A, Browne J, et al. The evidence on the effectiveness of management for malignant pleural effusion: a systematic review. *Eur J Cardiothorac Surg* 2006;29:829–838.
- Janssen JP, Collier G, Astoul P, et al. Safety of pleurodesis with talc poudrage in malignant pleural effusion: a prospective cohort study. *Lancet* 2007;369:1535–1539.
- Suzuki K, Servais EL, Rizk NP, et al. Palliation and pleurodesis in malignant pleural effusion: the role for tunneled pleural catheters. *J Thorac Oncol* 2011;6:762–767.
- Tremblay A, Michaud C. Single-center experience with 250 tunneled pleural catheter insertions for malignant pleural effusion. *Chest* 2006;129:362–368.
- Warren WH, Kalimi R, Khodadadian LM, et al. Management of malignant pleural effusions using the Pleur(x) catheter. *Ann Thorac Surg* 2008;85:1049–1055.
- Van Meter ME, McKee KY, Kohlwees RJ. Efficacy and safety of tunneled pleural catheters in adults with malignant pleural effusions: a systematic review. *J Gen Intern Med* 2011;26:70–76.
- Putnam JB Jr, Light RW, Rodriguez RM, et al. A randomized comparison of indwelling pleural catheter and doxycycline pleurodesis in the management of malignant pleural effusions. *Cancer* 1999;86:1992–1999.
- Demmy TL, Gu L, Burkhalter JE, et al. Optimal management of malignant pleural effusions (results of CALGB 30102). *J Natl Compr Canc Netw* 2012;10:975–982.
- Fysh ET, Waterer GW, Kendall PA, et al. Indwelling pleural catheters reduce inpatient days over pleurodesis for malignant pleural effusion. *Chest* 2012;142:394–400.
- Davies HE, Mishra EK, Kahan BC, et al. Effect of an indwelling pleural catheter vs chest tube and talc pleurodesis for relieving dyspnea in patients with malignant pleural effusion: the TIME2 randomized controlled trial. *JAMA* 2012;307:2383–2389.
- Freeman RK, Ascoti AJ, Mahidhara RS. A propensity-matched comparison of pleurodesis or tunneled pleural catheter in patients undergoing diagnostic thoracoscopy for malignancy. *Ann Thorac Surg* 2013;96:259–265.
- Hunt BM, Farivar AS, Vallières E, et al. Thoracoscopic talc versus tunneled pleural catheters for palliation of malignant pleural effusions. *Ann Thorac Surg* 2012;94:1053–1057.
- Puri V, Pyrdeck TL, Crabtree TD, et al. Treatment of malignant pleural effusion: a cost-effectiveness analysis. *Ann Thorac Surg* 2012;94:374–379.
- Refaat MM, Katz WE. Neoplastic pericardial effusion. *Clin Cardiol* 2011;34:593–598.
- Chong HH, Plotnick GD. Pericardial effusion and tamponade: evaluation, imaging modalities, and management. *Compr Ther* 1995;21:378–385.
- Wiener HC, Kristensen IB, Haubek A, et al. The diagnostic value of pericardial cytology: an analysis of 95 cases. *Acta Cytol* 1991;35:149–153.
- Clarke DP, Cosgrove DO. Real-time ultrasound scanning in the planning and guidance of pericardiocentesis. *Clin Radiol* 1987;38:119–122.
- Maher EA, Shepherd FA, Todd TJ. Pericardial sclerosis as the primary management of malignant pericardial effusion and cardiac tamponade. *J Thorac Cardiovasc Surg* 1996;112:637–643.
- Liu C, Crump M, Goss PE, et al. Prospective comparison of the sclerosing agents doxycycline and bleomycin for the primary management of malignant pericardial effusion and cardiac tamponade. *J Clin Oncol* 1996;14:3141–3147.
- Kunitoh H, Tamura T, Shibata T, et al. A randomised trial of intrapericardial bleomycin for malignant pericardial effusion with lung cancer (JCOG9811). *Br J Cancer* 2009;100:464–469.
- Ziskind AA, Pearce AC, Lemmon CC, et al. Percutaneous balloon pericardiectomy for the treatment of cardiac tamponade and large pericardial effusions: description of technique and report of the first 50 cases. *J Am Coll Cardiol* 1993;21:1–5.
- Allen KB, Faber LP, Warren WH, et al. Pericardial effusion: subxiphoid pericardiostomy versus percutaneous catheter drainage. *Ann Thorac Surg* 1999;67:437–440.
- Piehlner JM, Pluth JR, Schaff HV, et al. Surgical management of effusive pericardial disease: influence of extent of pericardial resection on clinical course. *J Thorac Cardiovasc Surg* 1985;90:506–516.
- Naunheim KS, Kesler KA, Fiore AC, et al. Pericardial drainage: subxiphoid vs. transthoracic approach. *Eur J Cardiothorac Surg* 1991;5:99–103, discussion 104.
- Vaitkus PT, Herrmann HC, LeWinter MM. Treatment of malignant pericardial effusion. *JAMA* 1994;272:59–64.
- Lestuzzi C. Neoplastic pericardial disease: Old and current strategies for diagnosis and management. *World J Cardiol* 2010;2(9):270–279.